



Honey-B-Healthy Use By Region


Management Survey 2012

Winter


Report ID: 185-2012

A comparison of average winter colony mortality among beekeepers who reported they did or did not apply and/or feed Honey-B-Healthy to a majority of their colonies, at least once, between April, 2011 and March, 2012 by region of operation.

No Significant Differences (within regions)



Respondent Ratio



Interpretation

No differences between groups.

Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April 1, 2011 and March, 2012?
-Honey B Healthy

		Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss		
				Mean	Standard Error	Mean(%)	Lower 95% CI	Upper 95% CI
Multiregional	Product Used	13	24,411	1877.8	1259.9	23.4	8.8	38.0
	Product Not Used	25	16,918	676.7	597.7	25.0	13.8	36.2
Northern States	Product Used	640	31,439	49.1	21.5	23.3	21.1	25.5
	Product Not Used	1,327	65,660	49.5	18.7	24.8	23.0	26.5
Southern States	Product Used	467	18,542	39.7	13.0	21.4	19.0	23.8
	Product Not Used	666	16,302	24.5	11.4	19.6	17.4	21.7

Comments About This Data

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.