



Carbohydrate Feed Use (Northeast)


Management Survey 2013

(Filtered by: Northeast)


A comparison of average winter colony mortality among beekeepers who did or did not feed carbohydrates to a majority of their colonies, at least once, between April and March.

Winter

Report ID: 164-2013



Participant Ratio



Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April 1, 2012 and March, 2013?

- Candy
- Dry sugar
- High Fructose Corn Syrup (HFCS)
- Honey
- Sugar syrup
- Wet supers

		Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss
				Mean	Standard Error	Mean(%) [Lower, Upper] CI
	Product Not Used	137	44,418	324.2	190.6	39.9 [34.5, 45.2]
	Product Used	958	50,249	52.5	22.6	49.3 [47.1, 51.6]

Comments About This Data

Relevant Links, References, and Citations



This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.