



# Single Stock (Northeast)


Management  
Survey 2013

( Filtered by:  
Northeast )

Average winter colony mortality reported by beekeepers who indicated which race of bees they kept between April and March.

## Winter

Report ID: 206-  
2013



## Survey Question

What type or race of bees do you use in your operation?

- New World Carniolan
- New World Carniolan hybrid
- Caucasian
- Caucasian hybrid
- Italian
- Italian hybrid
- Buckfast
- Buckfast hybrid
- Russian
- Russian hybrid
- Varroa Sensitive Hygienic (VSH)
- Minnesota Hygienic
- German black
- Local selected
- Africanized
- Don't know

		Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss
				Mean	Standard Error	Mean(%) [Lower, Upper] CI
Africanized	Not used at all or exclusively	590	14,355	24.3	9.1	52.5 [49.4,55.5]
Buckfast	Not used at all or exclusively	589	14,350	24.4	9.1	52.5 [49.5,55.6]
Buckfast hybrid	Not used at all or exclusively	585	14,339	24.5	9.2	52.6 [49.6,55.6]
	Used exclusively	5	16	3.2	0.7	36.7 [9.7,63.6]
Caucasian	Not used at all or exclusively	583	14,337	24.6	9.2	52.4 [49.4,55.4]
	Used exclusively	7	18	2.6	0.6	57.1 [24.8,89.5]
Caucasian hybrid	Not used at all or exclusively	587	14,313	24.4	9.2	52.6 [49.6,55.6]
Do Not Know	Not used at all or exclusively	973	53,151	54.6	22.0	49.0 [46.7,51.2]
	Used exclusively	136	14,477	106.4	99.2	45.1 [39.0,51.2]
German black	Not used at all or exclusively	589	14,349	24.4	9.1	52.4 [49.4,55.4]
Italian	Not used at all or exclusively	239	8,964	37.5	18.8	49.2 [44.3,54.1]
	Used exclusively	351	5,391	15.4	8.3	54.7 [50.9,58.5]
Italian hybrid	Not used at all or exclusively	546	13,966	25.6	9.8	52.3 [49.2,55.4]
	Used exclusively	44	389	8.8	3.0	54.9 [41.9,67.8]
Local selected	Not used at all or exclusively	528	8,218	15.6	6.0	54.7 [51.5,57.9]
	Used exclusively	62	6,137	99.0	69.8	33.4 [25.5,41.2]
Minnesota Hygienic	Not used at all or exclusively	588	14,348	24.4	9.1	52.5 [49.5,55.5]
New World Carniolan	Not used at all or exclusively	543	13,098	24.1	9.7	52.3 [49.1,55.4]
	Used exclusively	47	1,257	26.7	20.0	54.5 [43.5,65.5]
New World Carniolan hybrid	Not used at all or exclusively	575	14,299	24.9	9.3	52.4 [49.3,55.4]
	Used exclusively	15	56	3.7	0.7	55.3 [31.3,79.2]

Russian	Not used at all or exclusively	558	13,434	24.1	9.5	51.9 [48.8,55.0]
	Used exclusively	32	921	28.8	23.4	62.2 [47.7,76.6]
Russian hybrid	Not used at all or exclusively	590	14,355	24.3	9.1	52.5 [49.4,55.5]
Varroa Sensitive Hygienic (VSH)	Not used at all or exclusively	584	14,308	24.5	9.2	52.5 [49.5,55.6]
	Used exclusively	6	47	7.8	1.8	45.9 [12.9,78.8]
Single Queen Type Used	Only one breed used	590	14,355	24.3	9.1	52.5 [49.4,55.5]
	More then one type used	519	53,273	102.6	47.6	44.0 [41.1,46.9]

---

**Comments About This Data**

---

## Relevant Links, References, and Citations

Funded By:



United States  
Department of  
Agriculture

National Institute  
of Food  
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.