



Amitraz Use (East North Central)


Management Survey 2013

A comparison of average winter colony mortality among beekeepers who reported treating with Amitraz, not treating with a known varroa mite control product or not treating with an Amitraz based product (i.e. ApiVar), at least once, between April and March.


(Filtered by: East North Central)

Winter

Report ID: 273-2013



Participant Ratio



Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April 1, 2012 and March 2013?

-Amitraz based product

		Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss
				Mean	Standard Error	Mean(%) [Lower, Upper] CI
Amitraz	This Product Used	17	42,943	2526.1	836.2	29.9 [19.0, 40.9]
	Nothing Used	227	3,710	16.3	3.3	65.3 [60.9, 69.6]
	Other Product Used	390	37,433	96.0	38.6	59.7 [56.4, 63.0]

Comments About This Data

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.