



Formic Acid Use (Pennsylvania)


Management Survey 2013

(Filtered by: Pennsylvania)


Winter

Report ID: 277-2013

A comparison of average winter colony mortality among beekeepers who reported treating with formic acid, not treating with a known varroa mite control product, or not treating with a formic acid-based product, at least once, between April and March.



Participant Ratio



Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April, 2012 and March, 2013?

-Formic Acid

		Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss
				Mean	Standard Error	Mean(%) [Lower, Upper] CI
Formic	This Product Used	84	18,720	222.9	163.7	33.5 [27.6, 39.3]
	Nothing Used	186	6,298	33.9	22.7	60.1 [55.1, 65.1]
	Other Product Used	240	15,253	63.6	30.5	55.8 [51.2, 60.4]

Comments About This Data

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.