



Honey-B-Healthy Use By Region


Management Survey 2013

Winter


Report ID: 185-2013

A comparison of average winter colony mortality among beekeepers who reported they did or did not apply and/or feed Honey-B-Healthy to a majority of their colonies, at least once, between April and March.

No Significant Differences (within regions)



Respondent Ratio



Interpretation

No significant differences between the groups.

Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April 1, 2012 and March, 2013?

-Honey-B-Healthy

		Total Number of Respondents	Total Number of Colonies	Average Number of Colonies Managed	Average Colony Loss

		Providing Valid Responses	Managed			Mean(%) [Lower, Upper] CI
				Mean	Standard Error	
Multiregional	Product Used	41	236,708	5773.4	1874.6	29.5 [22.0,37.1]
	Product Not Used	87	192,716	2215.1	433.2	26.7 [21.5,31.8]
Northern States	Product Used	789	20,818	26.4	10.3	46.1 [43.7,48.6]
	Product Not Used	1,539	26,689	17.3	1.9	47.8 [46.0,49.6]
Southern States	Product Used	512	35,939	70.2	27.1	38.3 [35.4,41.2]
	Product Not Used	785	35,257	44.9	11.3	38.6 [36.2,41.1]

Comments About This Data

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.