




Coumaphos Use


A comparison of average winter colony mortality among beekeepers who reported treating with a Coumaphos based product, not treating with a known Varroa control product, or not treating with a Coumaphos-based product (e.g. CheckMite+), at least once, between April and March.

Winter

Report ID: 272-2013



Participant Ratio



Interpretation

Beekeepers who treated with Coumaphos did not report losing more or less colonies than those who did not report treating with Coumaphos.

Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April, 2012 and March, 2013?

-Coumaphos

| | | Total Number of Respondents Providing Valid Responses | Total Number of Colonies Managed | Average Number of Colonies Managed | | Average Colony Loss |
|-----------|--------------------|---|----------------------------------|------------------------------------|----------------|---------------------------|
| | | | | Mean | Standard Error | Mean(%) [Lower, Upper] CI |
| Coumaphos | This Product Used | 49 | 18,155 | 370.5 | 238.9 | 38.5 [29.1, 47.9] |
| | Nothing Used | 2,225 | 74,597 | 33.5 | 11.8 | 48.4 [46.9, 49.9] |
| | Other Product Used | 3,735 | 539,024 | 144.3 | 25.6 | 43.4 [42.3, 44.5] |

Comments About This Data

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.