




Fumagillin Use


A comparison of winter mortality among those who indicated they did or did not apply Fumagillin (registered for Nosema Control), at least once, to a majority of their colonies, between April and March.

Winter

Report ID: 33-2013



Participant Ratio



Interpretation

Beekeepers who treated with Fumagillin did not report losing more or less colonies than those who did not report treating with Fumagillin.

Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April, 2012 and March, 2013?

- Fluvinate- based product (Apistan)
- Amitraz- based product
- Coumaphos
- Amitraz
- Fumagillin

		Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss
				Mean	Standard Error	Mean(%) [Lower, Upper] CI
	Product Not Used	2,854	286,085	100.2	16.8	44.1 [42.8, 45.4]
	Product Used	930	271,094	291.5	89.7	41.1 [38.9, 43.2]

Comments About This Data

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.