




Fumagillin Use


A comparison of winter mortality among those who indicated they did or did not apply Fumagillin (registered for Nosema Control), at least once, to a majority of their colonies, between April 2010 and March 2011.

Winter

Report ID: 33



Respondent Ratio



Interpretation

No difference between groups

Survey Question

19. For the products listed below, indicate in which months you applied the product to a majority of your colonies.
Fumagillin

	Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss		
			Mean	Standard Error	Mean(%)	Lower 95% CI	Upper 95% CI
Product Used	902	115,404	127.9	26.8	34.4	32.1	36.7
Product Not Used	1,659	46,859	28.2	10.4	34.2	32.5	36.0

Comments About This Data

We do not know if the responding beekeepers applied this product according to label directions. It is likely that some of those who reported using this product did so using concentrations and delivery methods that differed from the label directions. This may affect the results reported here. More detailed questions regarding doses and application method are planned for future surveys.

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.