




MiteAThol Use By Region


A comparison of winter mortality among those who indicated they did or did not apply Mite-A-Thol (registered for the control of Honey Bee Tracheal Mites), at least once, to a majority of their colonies, between April 2010 and March 2011 by region of operation.

Winter

Report ID: 51



Respondent Ratio



Interpretation

No differences within groups

Survey Question

19. For the products listed below, indicate in which months you applied the product to a majority of your colonies.
Mite-A-Thol

		Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss		
				Mean	Standard Error	Mean(%)	Lower 95% CI	Upper 95% CI
Multiregional	Product Not Used	20	58,438	2921.9	877.0	22.4	13.6	31.3
Northern States	Product Used	56	509	9.1	3.4	45.4	35.5	55.3
	Product Not Used	1,235	15,021	12.2	1.3	40.5	38.4	42.6
Southern States	Product Used	47	434	9.2	1.6	18.7	11.5	25.8
	Product Not Used	851	19,623	23.1	5.2	25.5	23.4	27.7

Comments About This Data

Low use rate among responding beekeepers make comparisons difficult to interpret. Insufficient data to analyze use data by months

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.