



Drone Brood Removal Use (Northeast)



Management Survey 2013

(Filtered by: Northeast)

A comparison of average winter colony mortality among beekeepers who reported using or not using drone brood removal between April and March.

Winter

Report ID: 104-2013



Survey Question

Over the last year, did you employ any of the IPM practices/equipment listed below?

- Drone comb removal
- Small cell size comb
- Small hive beetle trap
- Screened bottom board

| | | Total Number of Respondents Providing Valid Responses | Total Number of Colonies Managed | Average Number of Colonies Managed | | Average Colony Loss |
|--|--------------------|---|----------------------------------|------------------------------------|----------------|---------------------------|
| | | | | Mean | Standard Error | Mean(%) [Lower, Upper] CI |
| | Technique Not Used | 657 | 6,056 | 9.2 | 1.1 | 51.1 [48.4, 53.8] |
| | Technique Used | 213 | 8,998 | 42.2 | 21.0 | 41.7 [37.2, 46.3] |

Comments About This Data

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.