
INFORMED.

Antibiotic Use By Product

Winter

Report ID: 60-2014 A comparison of average winter colony mortality among beekeepers who applied or did not apply either Terramycin or Tylosin (Tylan), at least once, to a majority of their colonies between April and March.



Interpretation

Beekeepers who treated with Tylan antibiotics reporting losing 9 fewer overwintering colony deaths per 100 managed colonies than those who did not report treating with antibiotics. In other words, beekeepers who reported treating with Tylan antibiotics lost 20.7% fewer colonies than the ones who did not report treating with any known antibiotic product. There is no significant difference between the Terramycin groups.

Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April, 2013 and March, 2014?

- -Terramycin
- -Tylosin (Tylan)

		Respondents	Number of	Average Number of Colonies Managed		Average Colony Loss
				Mean	Standard Error	Mean(%) [Lower, Upper] CI
Terramycin	Product Used	400	322 , 334	805.8	204.4	40.1 [36.8,43.3]

	Product Not Used	5 , 221	204,255	39.1	6.5	43.6 [42.6,44.6]
Tylan	Product Used	110	333 , 154	3028.7	720.2	34.5 [29.2,39.9]
	Product Not Used	5 , 511	193,435	35.1	5.3	43.5 [42.5,44.5]

Comments About This Data

Relevant Links, References, and Citations

Funded By:





United States Department of Agriculture National Institute of Food and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.