




Fluvalinate Use


A comparison of average winter colony mortality among beekeepers who reported treating with Fluvalinate, not treating with any known Varroa control product, or not treating with a Fluvalinate-based product (e.g. Apistan), at least once, between April and March.

Winter

Report ID: 271-2014



Participant Ratio



Interpretation

Beekeepers who reported treating with Fluvalinate reported 15.2 fewer overwintering colony deaths per 100 managed colonies than those who did not report using a known Varroa mite control product. In other words, beekeepers who reported treating with Fluvalinate lost 31.1% fewer colonies than those who did not report treating with any known Varroa mite control product.

Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April, 2013 and March, 2014?

-Fluvalinate

		Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss
				Mean	Standard Error	Mean(%) [Lower, Upper] CI
Fluvalinate	This Product Used	121	14,924	123.3	50.0	33.7 [27.7, 39.6]
	Nothing Used	3,269	99,042	30.3	20.9	48.9 [47.5, 50.2]
	Other Product	2,231	412,623	184.9	25.8	35.8 [34.4, 37.2]

Used

Comments About This Data

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.