




Honey-B-Healthy Use By Region

Management Survey 2014


Winter

Report ID: 185-2014

A comparison of average winter colony mortality among beekeepers who reported they did or did not apply and/or feed Honey-B-Healthy to a majority of their colonies, at least once, between April and March.



Respondent Ratio



Interpretation

No significant differences between the groups.

Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April 1, 2013 and March, 2014?

-Honey-B-Healthy

		Total Number of Respondents	Total Number of Colonies	Average Number of Colonies Managed	Average Colony Loss

		Providing Valid Responses	Managed			
				Mean	Standard Error	Mean(%) [Lower, Upper] CI
Multiregional	Product Used	49	226,070	4613.7	1445.8	21.1 [15.6, 26.6]
	Product Not Used	57	180,452	3165.8	591.1	28.2 [21.0, 35.3]
Northern States	Product Used	1,119	15,681	14.0	1.9	46.9 [44.7, 49.1]
	Product Not Used	2,550	27,879	10.9	1.8	50.2 [48.7, 51.7]
Southern States	Product Used	643	31,323	48.7	19.3	32.1 [29.5, 34.6]
	Product Not Used	1,146	38,093	33.2	13.7	34.0 [32.0, 36.1]

Comments About This Data

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.