





A comparison of winter mortality among those who indicated they applied or did not apply Nosevet, at least once, to a majority of their colonies between April and March.

Winter

Report ID: 243-2014



Interpretation

Beekeepers who treated with Nosevet did not report losing more or less colonies than those who did not report treating with Nosevet.

Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April 1, 2013 and March, 2014?

-Nosevet

		Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss
				Mean	Standard Error	Mean(%) [Lower, Upper] CI
Nosevet	Product Not Used	5,518	488,095	88.5	16.1	43.3 [42.3, 44.3]
	Product Used	103	38,494	373.7	128.3	43.2 [36.6, 49.9]

Relevant Links, References, and Citations

Funded By:



United States
Department of
Agriculture

National Institute
of Food
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.