





Average winter colony mortality reported by beekeepers who reported they reused or did not reuse brood comb.

Winter

Report ID: 71-2015



Participant Ratio



Interpretation

Beekeepers who reported reusing brood comb did not suffer fewer or greater losses than beekeepers who did not reuse any old brood comb.

Survey Question

Between April, 2014 and March, 2015, before you re-used brood comb that you had taken out of production or purchased, did you...

Check all that apply

- I did not reuse any old brood comb
- Cull any particularly old or bad combs and replace them
- Irradiate the comb
- Fumigate the comb with acetic acid
- Freeze the comb
- Store the comb with paradichlorobenzene crystals
- Did not treat the comb in any particular way
- Other (please specify)

		Total Number of Respondents Providing Valid Responses	Total Number of Colonies Managed	Average Number of Colonies Managed		Average Colony Loss
				Mean	Standard Error	Mean(%) [Lower, Upper] CI
Disposed Used Brood Comb	Combs Not Reused	1,211	25,024	20.7	9.4	40.0 [37.8, 42.3]

Combs Reused	3,162	312,528	98.8	24.1	42.7 [41.5, 43.9]
--------------	-------	---------	------	------	-------------------

Comments About This Data

Relevant Links, References, and Citations



This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.