



# Honey-B-Healthy Use By Region


Management Survey 2015

## Winter


Report ID: 185-2015

A comparison of average winter colony mortality among beekeepers who reported they did or did not apply and/or feed Honey-B-Healthy to a majority of their colonies, at least once, between April and March.

No Significant Differences (within regions)



## Respondent Ratio



## Interpretation

There are no significant differences within regions.

## Survey Question

Which, if any of the following, did you apply to a majority of your colonies between April 1, 2014 and March, 2015?

-Honey-B-Healthy

		Total Number of Respondents	Total Number of Colonies Managed	Average Number of Colonies Managed	Average Colony Loss

		Providing Valid Responses				Mean(%) [Lower, Upper] CI
				Mean	Standard Error	
Multiregional	Product Used	37	129,786	3507.7	1733.4	36.7 [27.6, 45.9]
	Product Not Used	113	124,073	1098.0	273.1	26.7 [21.7, 31.7]
Northern States	Product Used	956	22,307	23.3	10.2	43.9 [41.7, 46.2]
	Product Not Used	2,016	26,662	13.2	2.7	47.4 [45.8, 49.0]
Southern States	Product Used	582	13,990	24.0	7.8	36.0 [33.2, 38.7]
	Product Not Used	927	43,092	46.5	21.3	35.2 [33.0, 37.4]

## Comments About This Data

## Relevant Links, References, and Citations

Funded By:



United States  
Department of  
Agriculture

National Institute  
of Food  
and Agriculture

This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by the Bee Informed Partnership or its members. The results presented here are the summary of the population who responded. The sample may not be representative of the beekeeping population at large. These results simply highlight differences in the sample population. The results cannot be considered conclusive, causative, protective, or attest to product efficacy or lack of efficacy.